

According to the Noether theorem, a translationally invariant system of classical fields  $\phi_a$  has a conserved stress-energy tensor

$$T_{\text{Noether}}^{\mu\nu} = \sum_a \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi_a)} \partial^\nu \phi_a - g^{\mu\nu} \mathcal{L}. \quad (1)$$

Actually, to assure the symmetry of the stress-energy tensor,  $T^{\mu\nu} = T^{\nu\mu}$  (which is necessary for the angular momentum conservation), one sometimes has to add a total divergence,

$$T^{\mu\nu} = T_{\text{Noether}}^{\mu\nu} + \partial_\lambda \mathcal{K}^{[\lambda\mu]\nu}, \quad (2)$$

where  $\mathcal{K}^{[\lambda\mu]\nu}$  is some 3-index Lorentz tensor antisymmetric in its first two indices.

1. Show that regardless of the specific form of  $\mathcal{K}^{[\lambda\mu]\nu}(\phi, \partial\phi)$ ,

$$\begin{aligned} \partial_\mu T^{\mu\nu} &= \partial_\mu T_{\text{Noether}}^{\mu\nu} = (\text{hopefully}) = 0 \\ P_{\text{net}}^\mu &\equiv \int d^3\mathbf{x} T^{0\mu} = \int d^3\mathbf{x} T_{\text{Noether}}^{0\mu}. \end{aligned} \quad (3)$$

For the scalar fields, real or complex,  $T_{\text{Noether}}^{\mu\nu}$  is properly symmetric and one simply has  $T^{\mu\nu} = T_{\text{Noether}}^{\mu\nu}$ . Unfortunately, the situation is more complicated for the vector, tensor or spinor fields. To illustrate the problem, consider the free electromagnetic fields described by the Lagrangian

$$\mathcal{L}(A_\mu, \partial_\nu A_\mu) = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} \quad (4)$$

where  $A_\mu$  is a real vector field and  $F_{\mu\nu} \equiv \partial_\mu A_\nu - \partial_\nu A_\mu$ .

2. Write down  $T_{\text{Noether}}^{\mu\nu}$  for the free electromagnetic fields and show that it is neither symmetric nor gauge invariant.
3. The properly symmetric — and also gauge invariant — stress-energy tensor for the free electromagnetism is

$$T_{\text{EM}}^{\mu\nu} = -F^{\mu\lambda} F^\nu{}_\lambda + \frac{1}{4} g^{\mu\nu} F_{\kappa\lambda} F^{\kappa\lambda}. \quad (5)$$

Show that this expression indeed has form (2) for some  $\mathcal{K}^{[\lambda\mu]\nu}$ .

4. Write down the components of the stress-energy tensor (5) in non-relativistic notations and make sure you have the familiar electromagnetic energy density, momentum density and pressure.

Now consider the electromagnetic fields coupled to a conserved electric current  $J^\mu$ .

5. Use Maxwell's equations to show that

$$\partial_\mu T_{\text{EM}}^{\mu\nu} = -F^{\nu\lambda} J_\lambda. \quad (6)$$

Eq. (6) suggests that any system of charged “matter” fields which carries the current  $J^\mu$  should have its stress-energy tensor  $T_{\text{mat}}^{\mu\nu}$  obeying

$$\partial_\mu T_{\text{mat}}^{\mu\nu} = +F^\nu_\lambda J_{\text{EM}}^\lambda. \quad (7)$$

Consequently, the combined stress-energy tensor  $T_{\text{EM}}^{\mu\nu} + T_{\text{mat}}^{\mu\nu}$  should be divergence-less and thus lead to a conserved total energy and momentum.

Generally, testing eq. (7) for any particular system of charged “matter” fields makes use of fields' equations of ‘motion’ and also of the fact that *the covariant derivatives  $D_\mu$  do not commute with each other*. Instead, when acting upon a field  $\Phi_q$  of charge  $q$ , one has

$$(D_\mu D_\nu - D_\nu D_\mu)\Phi_q = iq F_{\mu\nu} \Phi_q \quad (8)$$

(in  $c = \hbar = 1$  units).

6. Verify eq. (8).

Consider a specific example of the EM coupled to a complex scalar field of electric charge  $q \neq 0$ . The net Lagrangian density of the EM + scalar fields is

$$\mathcal{L} = D^\mu \Phi^* D_\mu \Phi - m^2 \Phi^* \Phi - \frac{1}{4} F^{\mu\nu} F_{\mu\nu}. \quad (9)$$

7. Verify the gauge invariance of the electric current

$$J^\mu = -\frac{\partial \mathcal{L}}{\partial A_\mu} \quad (10)$$

by expressing it in terms of the scalar fields and their *covariant* derivatives  $D^\mu \Phi$  and  $D^\mu \Phi^*$ .

8. Write down equations of motion for the scalar fields  $\Phi$  and  $\Phi^*$  and use them to verify the current conservation  $\partial_\mu J^\mu = 0$ .
9. Write down the Noether stress-energy tensor for this field system and show that

$$T_{\text{net}}^{\mu\nu} \equiv T_{\text{EM}}^{\mu\nu} + T_{\text{mat}}^{\mu\nu} = T_{\text{Noether}}^{\mu\nu} + \partial_\lambda \mathcal{K}^{[\lambda\mu]\nu} \quad (11)$$

where  $\mathcal{K}^{[\lambda\mu]\nu}$  is the same function of EM fields as in the free EM case [3.],  $T_{\text{EM}}^{\mu\nu}$  is exactly as in eq. (5), and

$$T_{\text{mat}}^{\mu\nu} = D^\mu \Phi^* D^\nu \Phi + D^\nu \Phi^* D^\mu \Phi - g^{\mu\nu} (D_\lambda \Phi^* D^\lambda \Phi - m^2 \Phi^* \Phi) \quad (12)$$

Hint: In the presence of an electric current  $J^\mu$ , the  $\partial_\lambda \mathcal{K}^{[\lambda\mu]\nu}$  correction to the electromagnetic stress-energy tensor contains an extra  $J^\mu A^\nu$  term. This term is important for obtaining a gauge-invariant stress-energy tensor (12) for the scalar field.

10. Use the scalar field's equations of motion and eq. (8) to verify eq. (7).